

**Corrigendum to Yasunaga T, Chérot F & Schwartz MD (2021) New genera and species of the Oriental mirine plant bugs from Southeast Asia, with six new combinations (Insecta: Heteroptera: Miridae: Mirinae: Mirini). Raffles Bulletin of Zoology, 69: 137–155.**

P. 142, Table 1. The genus name for the two new species, *Sundacapsus flavonitidus* and *Sundacapsus kinabaluensis*, was incorrectly shown as *Sumatracapsus*. The full table with the corrected genus names is provided below.

Table 1. Measurements for Oriental mirine species. Abbreviations: L = length, W = width.

	Body	Head	Vertex	Pronotum		Max	Antennomere L				Labium	Hind leg segments L		
	L	W	W	L	W	W	I	II	III	IV	L	Femur	Tibia	Tarsus
<i>Malayamiridius nigrotenuis</i> male (n = 3)														
max	5.13	0.95	0.29	1.03	1.38	1.41	0.95	2.34	2.09	0.93	2.22	2.28	3.61	0.55
min	4.80	0.93	0.23	0.95	1.32	1.35	0.87	2.13	1.84	0.82	1.90	2.09	3.23	0.51
mean	<b>4.98</b>	<b>0.94</b>	<b>0.25</b>	<b>0.98</b>	<b>1.36</b>	<b>1.37</b>	<b>0.92</b>	<b>2.22</b>	<b>1.94</b>	<b>0.87</b>	<b>2.01</b>	<b>2.17</b>	<b>3.41</b>	<b>0.53</b>
<i>Malayamiridius nigrotenuis</i> female (n = 3)														
max	6.06	0.99	0.34	1.15	1.65	1.71	0.99	2.57	2.30	0.95	2.24	2.57	3.99	0.59
min	5.49	0.97	0.30	1.03	1.58	1.58	0.84	2.32	1.98	0.93	1.82	2.28	3.57	0.55
mean	<b>5.78</b>	<b>0.98</b>	<b>0.33</b>	<b>1.11</b>	<b>1.62</b>	<b>1.63</b>	<b>0.93</b>	<b>2.46</b>	<b>2.09</b>	<b>0.94</b>	<b>2.08</b>	<b>2.44</b>	<b>3.82</b>	<b>0.57</b>
<i>Sundacapsus flavonitidus</i> male (n = 4)														
max	5.88	1.08	0.35	1.24	1.79	2.01	1.14	2.89	1.73	0.61	2.09	2.34	3.90	0.76
min	4.86	0.99	0.32	1.12	1.69	1.79	1.01	2.66	1.56	0.57	2.05	2.15	3.33	0.68
mean	<b>5.33</b>	<b>1.03</b>	<b>0.34</b>	<b>1.15</b>	<b>1.72</b>	<b>1.88</b>	<b>1.09</b>	<b>2.79</b>	<b>1.66</b>	<b>0.59</b>	<b>2.03</b>	<b>2.22</b>	<b>3.60</b>	<b>0.73</b>
<i>Sundacapsus flavonitidus</i> female (n = 3)														
max	6.48	1.14	0.40	1.35	2.09	2.32	0.95	2.32	1.52	0.61	2.20	2.34	3.25	0.74
min	5.76	1.06	0.38	1.31	1.92	2.20	0.82	2.28	1.39	0.59	2.05	2.26	3.19	0.70
mean	<b>6.09</b>	<b>1.10</b>	<b>0.39</b>	<b>1.32</b>	<b>2.02</b>	<b>2.26</b>	<b>0.87</b>	<b>2.30</b>	<b>1.45</b>	<b>0.60</b>	<b>2.12</b>	<b>2.29</b>	<b>3.22</b>	<b>0.72</b>
<i>Sundacapsus kinabaluensis</i> male (n = 4)														
max	7.10	1.14	0.34	1.28	2.13	2.30	1.23	2.85	1.28	0.65	2.40	2.75	3.98	0.88
min	5.90	1.04	0.30	1.13	1.88	2.04	1.00	2.44	1.23	0.60	2.18	2.33	3.50	0.85
mean	<b>6.58</b>	<b>1.09</b>	<b>0.32</b>	<b>1.20</b>	<b>1.99</b>	<b>2.17</b>	<b>1.10</b>	<b>2.60</b>	<b>1.26</b>	<b>0.63</b>	<b>2.28</b>	<b>2.49</b>	<b>3.68</b>	<b>0.87</b>
<i>Sundacapsus kinabaluensis</i> female (n = 5)														
max	7.93	1.20	0.43	1.50	2.54	2.80	1.00	2.34	1.30	0.68	2.48	2.75	4.10	0.88
min	7.60	1.15	0.39	1.43	2.48	2.65	0.98	2.10	1.16	0.63	2.36	2.50	3.70	0.80
mean	<b>7.78</b>	<b>1.17</b>	<b>0.40</b>	<b>1.46</b>	<b>2.50</b>	<b>2.72</b>	<b>0.99</b>	<b>2.21</b>	<b>1.23</b>	<b>0.65</b>	<b>2.42</b>	<b>2.59</b>	<b>3.80</b>	<b>0.82</b>
<i>Tondomiris puguis</i> male														
Holotype	<b>6.30</b>	<b>1.08</b>	<b>0.32</b>	<b>1.25</b>	<b>2.01</b>	<b>2.28</b>	<b>1.25</b>	<b>2.62</b>	<b>2.09</b>	–	<b>3.10</b>	<b>3.14</b>	<b>4.66</b>	<b>0.76</b>