

Biodiversity Record: The mangrove crab, *Neosarmatium smithi*, on Saint John's Island

Bee Yan Lee*, Yu Xun Leow & Hashani Dharan

Tropical Marine Science Institute, National University of Singapore, Singapore 119227

Email: tmslby@nus.edu.sg (*corresponding author)

Recommended citation. Lee BY, Leow YX & Dharan H (2025) Biodiversity Record: The mangrove crab, *Neosarmatium smithi*, on Saint John's Island. *Nature in Singapore*, 18: e2025052. DOI: 10.26107/NIS-2025-0052

Subject: *Neosarmatium smithi* (Malacostraca: Decapoda: Brachyura: Sesarmidae)

Subject identified by: Bee Yan Lee.

Location, date and time: Singapore Strait, Saint John's Island; 13 January 2025; around 0835 hrs

Habitat: Parkland, on a concrete path about 300 m from a patch of mangrove forest (Bendera Bay mangrove).

Observers: Hashani Dharan and Leow Yu Xun.

Observation: The subject, around 30 mm in carapace width, was found crouching motionless on wet concrete ground, on a rainy day, outside the Saint John's Island National Marine Laboratory.




Fig. 1. In-situ dorso-frontal view of *Neosarmatium smithi*. (Photograph by: Leow Yu Xun). Fig. 2. Frontal view of the same animal (Photograph by: Hashani Dharan).

Remarks: *Neosarmatium smithi* is classified as an obligate mangrove dweller in Singapore (Tan & Ng, 1994) where it was hitherto known only from the northern part of Singapore, in patches of mangroves along the Johor Strait. Based on holdings of this species in the Lee Kong Chian Natural History Museum at the National University of Singapore, as well as field observations by the first author, *Neosarmatium smithi* has been recorded from Lim Chu Kang, Mandai, Loyang, Pasir Ris, and Pulau Tekong. The present sighting marks the first record for this species in the southern part of the country in the Singapore Strait.

Neosarmatium smithi is distributed in the Indo-West Pacific region, from southern Africa and Madagascar eastwards to the Philippines, China, and Japan (Davie, 1994). The species typically inhabits mangrove forests where it feeds on fallen leaves (Davie, 1994; Fratini et al., 2000). By consuming leaf litter, it may play an important role in the nutrient cycle by facilitating the reduction of iron compounds in the environment (Nakanishi et al., 2020).

Literature cited:

Davie PJF (1994) Revision of *Neosarmatium* Serène and Soh (Crustacea: Brachyura: Sesarminae) with description of two new species. *Memoirs of the Queensland Museum*, 35(1): 35–74.

Fratini S, Cannicci S & Vannini M (2000) Competition and interaction between *Neosarmatium smithi* (Crustacea: Grapsidae) and *Terebralia palustris* (Mollusca: Gastropoda) in a Kenyan mangrove. *Marine Biology*, 137: 309–316.

Nakanishi Y, Matsutani T, Hinokidani K, Nagai T & Irie M (2020) A role of a herbivorous crab, *Neosarmatium smithi*, in dissolved iron elution from mangrove ecosystems. *Tropics*, 28(4): 91–97.

Tan CGS & Ng PKL (1994) An annotated checklist of mangrove brachyuran crabs from Malaysia and Singapore. *Hydrobiologia*, 28: 75–84.